ON COMPOSITION OPERATORS ON A^2

BY
MAHMUD ILAYYAN MASRI
DEPARTMENT OF MATHEMATICS
AN-NAJAH NATIONAL UNIVERSITY

ABSTRACT

If Φ is an analytic function mapping the open unit disk D into itself and A^2 is the Bergman space of analytic functions on D, the compositon operator C_Φ on A^2 is defined by $C_\phi f = fo\Phi \forall f \varepsilon A^2$.

In this paper we consider the spectral radius, unitary equivalence, subnormality of C_Φ and study the case $\Phi(z) = z^m$, m = 2, 3, ... in detail.

ملخص

اذا كانت Φ داله تحليليه من القرص المفتوح D الى نفسه و A^2 هو فراغ برغمان المكون من الدوال التحليليه على D والتي تحقق شرط تكامل معين فانه يمكن تعريف المؤثر المركب C_Φ على A^2 كما يلي: $C_\phi f = f \circ \phi$ لكل f تنتمي الى A^2.

في ورقة البحث هذه ندرس نصف القطر الطيفي والتكافؤ الأحادي والصفه شبه الطبيعيه للمؤثر المذكور أعلاه. وكذلك نبحث بشيء من التفصيل في الحاله الخاصه
$\Phi(z) = z^m$, m = 2, 3, ...

Hence, the norm of f is given by

$$\|f\|^2 = \langle f, f\rangle = \sum_{n=0}^{\infty} \frac{|a_n|^2}{n+1}$$

Of special interest is the function $K_\zeta(z) = (1-\zeta z)^{-2}$ which serves as the "reproducing kernel" for A^2, i.e.,

$$f(\zeta) = \langle f, k_\zeta\rangle \forall\ f \in A^2 \ \&\ \forall \zeta \in D.$$

Furthermore, the functions $e_n(z) = \sqrt{n+1}\ z^n$, $n = 0,1,2,\ldots$ form an orthonormal basis for A^2.

If Φ is a non-constant analytic function mapping D into itself, then Φ induces a composition operator $C_\Phi : A^2 \to A^2$ defined by $C_\Phi f = f\circ\Phi\ \forall\ f \in A^2$.

Boyd [1] showed that C_Φ is bounded and obtained norm estimats for C_Φ. He studied normal, unitary, hermetian and compact composition operators on A^2. Furthermore, he computed the spectrum of C_Φ for some special kinds of Φ.

Cowen [6] computed the spectral radius of C_Φ as an operator on the Hardy space H^2. Here, we compute the spectral radius of C_Φ as an operator on A^2. Also, in recent work Campbell-Wright [2] found a necessary and sufficient condition for two composition operators on H^2 to be unitarily equivalent.We show that the same thing holds in the case of A^2. Moreover, we give a necessary condition for the subnormality of C_Φ on A^2.

Finally, as an example we study C_Φ when $\Phi(z) = z^m$, $m = 2,3,\ldots$

2.Spectral raduis. It was found out that the fixed points of Φ are related to some properities of C_Φ and to its spectral radius in particular. We say that a point $b \in \overline{D}$, the closure of D, is a fixed point of Φ if $\text{limit}_{r\to 1^-}\Phi(rb) = b$. We write $\text{limit}_{r\to 1^-}\Phi'(rb) = \Phi'(b)$. Although it is not apriori evident that Φ has fixed points the following is known.

Denjoy-Wolff Theorem [8,9] : let $\Phi : D \to D$ be analytic and non-elliptic Mobius transformation onto D. Then $\exists$ a unique fixed point a of Φ in $\overline{D}$ such that $|\Phi'(a)| \leq 1$.

We call the distinguished fixed point a the Denjoy-Wolff point of Φ and we point out that if $|a| = 1$, then $0 < \Phi'(a) \leq 1$ and if $|a|<1$, then $0 \leq |\Phi'(a)| < 1$. Now we are ready to prove the spectral radius theorem which is similar to that of [6] in the H^2 case.

Spectral radius theorem : Let $\Phi : D \to D$ be analytic with Denjoy-Wolff Point a. Then the spectral radius $r(C_\Phi)$ of C_Φ is 1 when $|a| < 1$ and $(\Phi'(a))^{-1}$ when $|a| = 1$.

Proof : $r(C_\Phi) = \text{limit}_{n\to\infty} \| C^n_\Phi \|^{1/n}$

$= \text{limit}_{n\to\infty} \| C_{\Phi^n} \|^{1/n}$

Where $\Phi_n = \Phi 0 \Phi_{n-1}$, $n=1,2,\ldots$ $\Phi_1 = \Phi$ and $\Phi_0(z) = z$ $\forall$ $z \in D$ (see e.g., [4,p.142]).

Boyd [1] showed that

$$(1-|\phi(0)|^2)^{-1} \leq \|C_\phi\| \leq \frac{1+|\phi(0)|}{1-|\phi(0)|}$$

Hence,

$$\lim_{n\to\infty} sup\ (1+|\phi_n(0)|^2)^{-1/n} \leq r(C_\phi) \leq$$

$$\lim_{n\to\infty} inf\ \left(\frac{1+|\phi_n(0)|}{1-|\phi_n(0)|}\right)^{1/n}$$

Since,

$$\lim_{n\to\infty} inf\ \left(\frac{1+|\phi_n(0)|}{1-|\phi_n(0)|}\right)^{1/n} =$$

$$\lim_{n\to\infty} inf\ (1+|\phi_n(0)|)^{2/n}(1-|\phi_n(0)|^2)^{-1/n} \cdot$$

$$= \lim_{n\to\infty} inf\ (1-|\phi_n(0)|^2)^{-1/n}$$

We have

$$r(C_\phi) = limit_{n\to\infty}(1-|\phi_n(0)|^2)^{-1/n}$$

$$= limit_{n\to\infty}(1-|\phi_n(0)|)^{-1/n} \qquad (2.1)$$

Since (see [5]) $\lim_{n\to\infty} \Phi_n(0) = a$, $r(C_\Phi) = 1$ if $|a| < 1$ by (2.1). When $|a| = 1$ and $\Phi'(a) < 1$ we have [3,p.32]

$$limit_{n\to\infty}\frac{1-|\phi_n(0)|}{1-|\phi_{n-1}(0)|} = \phi'(a)$$

Therefore, (2.1) implies

$$r(C_\phi) = limit_{n\to\infty}(1-|\phi_n(0)|)^{-1/n}$$

$$= limit_{n\to\infty}\left(\prod_{k=0}^{n-1}\frac{1-|\phi_k(0)|}{1-|\phi_{k+1}(0)|}\right)^{1/n}$$

$$= limit_{n\to\infty}\frac{1-|\phi_{n-1}(0)|}{1-|\phi_n(0)|} = (\phi'(a))^{-1}$$

Next, suppose that $|a| = 1$ and $\Phi'(a) = 1$. If $\{z_n\}$ is a sequence in D converging to a such that $\Phi(z_n) \to a$ as $n \to \infty$ and

$$\alpha = limit_{n\to\infty} \frac{1-|\phi(z_n)|}{1-|z_n|}$$

exists then by [3,pp25-32] $\alpha \geq \Phi'(a) = 1$. Hence , letting
$z_n = \Phi_{n-1}(0)$, $n = 1,2, \ldots$, we get

$$\lim_{n\to\infty} inf \ \frac{1-|\phi_n(0)|}{1-|\phi_{n-1}(0)|} \geq 1.$$

Therefore, by (2.1)

$$r(C_\phi) = limit_{n\to\infty} (\Pi_{k=0}^{n-1} \frac{1-|\phi_k(0)|}{1-|\phi_{k+1}(0)|})^{1/n}$$

$$\leq \lim_{n\to\infty} sup \ \frac{1-|\phi_{n-1}(0)|}{1-|\phi_n(0)|} \leq 1$$

But (2.1) again implies $r(C_\Phi) \geq 1$ since $1-|\Phi_n(0)| \leq 1$ $\forall$ n = 1,2,... . Thus $r(C_\Phi) = 1 = (\Phi'(a))^{-1}$.

3. Unitary equivalence. Campbell-Wright [2] proved a theorem concerning unitary equivalence of composition operators on H^2. Here we show that the same thing holds in the A^2 case.

Theorem : Let Φ and Ψ be analytic functions, not disk automorphisms, that map D into itself. Suppose that the Denjoy-Wolff point a of Φ is in D with $\Phi_n(0) \neq a$ $\forall$ positive integers n . Then C_Φ is unitarily equivalent to C_Ψ on A^2 iff $\Psi(z) = e^{i\theta}\Phi(e^{-i\theta}z)$ for some real number θ.

Proof : Let U be a unitary operator on A^2 such that $C_\Psi = U^* C_\Phi U$. Since $|a| < 1$ we have $0 \leq |\Phi'(a)| < 1$. Thus [5] implies that the non-zero solutions of the equation $f\circ\Phi = f$ are the constant functions. Hence, the same is true for the equation $f\circ\Psi = f$ by the unitary equivalence of C_Φ and C_Ψ. Therefore, $U(1) = \gamma 1$ where $|\gamma| = 1$. Since $K_0 = 1$ and $C^*_\Phi k_\alpha = K_{\Phi(\alpha)}$ where C^*_Φ is the adjoint of C_Φ, it follows that $\forall$ n

$$UK_{\Psi_n(0)} = UC^*_{\Psi_n}(k_0) = UC^{*n}_{\Psi}(k_0) = C^{*n}_{\Phi}U(K_0) = \gamma C^{*n}_{\Phi}(k_0) = \gamma K_{\Phi_n(0)}$$

In particular, when n = 1, we get

$$(1-|\psi(0)|^2)^{-1}=\|k_{\psi(0)}\|=\|Uk_{\psi(0)}\|=\|\gamma k_{\phi(0)}\|$$

$$=\|k_{\phi(0)}\|=(1-|\phi(0)|^2)^{-1}$$

Therefore, $\Phi(0) = e^{-i\theta}\,\Psi(0)$ for some real number θ. Furthermore,

$$(1-\overline{\Phi}(0)\phi_n(0))^{-2}=k_{\phi(0)}(\phi_n(0))=\langle k_{\phi(0)}, k_{\phi_n(0)}\rangle$$

$$=\langle \gamma k_{\phi(0)}, \gamma k_{\psi_n(0)}\rangle=\langle Uk_{\psi(0)}, Uk_{\psi_n(0)}\rangle$$

$$=\langle k_{\psi(0)}, k_{\psi_n(0)}\rangle = k_{\psi(0)}(\psi_n(0)$$

$$=(1-\overline{\psi}(0)\psi_n(0))^{-2}$$

Thus, $\Phi_n(0) = (\overline{\Psi(0)}/\overline{\Phi(0)})\,\Psi_n(0) = e^{-i\theta}\,\Psi_n(0)$. It follows that the analytic functions $\Psi(z)$ and $e^{i\theta}\,\Phi(e^{-i\theta}z)$ agree on the sequence $\{e^{i\theta}\,\Phi_n(0)\}$ which converges to $e^{i\theta}$ a in D and hence $\Psi(z) = e^{i\theta}\Phi(e^{-i\theta}z)$.

Conversely, if $\beta(z) = e^{i\theta}\,z$, $z \in D$, then by [1] C_β is a unitary operator on A^2 and $C_\Psi = C^*_\beta C_\Phi C_\beta$.

4. Subnormality of C_Φ on A^2 . Boyd [1] proved that C_Φ is normal on A^2 iff $\Phi(z) = \alpha z$ for some α with $|\alpha| \le 1$ iff C^*_Φ is a composition operator. Here, we give a necessary condition for the subnormality of C_Φ on A^2. Let S be an operator on a Hilbert space

H. S is called subnormal if there is a Hilbert space K containing H and a normal operator N on K such that N leaves H invariant and S is the restriction of N to H. Also, S is called hyponormal if $S^* S \geq S S^*$ where S^* is the adjoint of S.

Theorem 4.1 : If $\exists$ a positive integer n such that

$$\|C_\phi^2 e_n\| < \|C_\phi e_n\|^2 \qquad (4.1)$$

then C_Φ is not subnormal.

Proof : Suppose $\exists$ n as in (4.1). Let $f_o = \beta e_n$ and $f_1 = \gamma e_n$ where β and $\gamma \in \mathbb{R}$. It follows that

$$\sum_{j,k=0}^{1} \langle C_\phi^{j+k} f_j, C_\phi^{j+k} e_k \rangle -$$

$$\langle f_o, f_o \rangle + \langle C_\phi f_1, C_\phi f_o \rangle + \langle C_\phi f_o, C_\phi f_1 \rangle + \langle C_\phi^2 f_1, C_\phi^2 f_1 \rangle$$

$$-\beta^2 + 2\beta\gamma \|C_\phi e_n\|^2 + \gamma^2 \|C_\phi^2 e_n\|$$

$$-g(\beta, \gamma)$$

Hence, (4.1) implies that the function $g(\beta,\gamma)$ has a saddle point at $(0,0)$. Thus $\exists$ non-zero $\beta,\gamma \in \mathbb{R}$ such that

$$\sum_{j,k=0}^{1} \langle C_{\Phi}^{k+j} f_j, C_{\Phi}^{j+k} f_k \rangle < 0.$$

Therefore, [4,p.117] implies that C_{Φ} is not subnormal.

In [7] Cowen and Kriete proved that $\Phi(0) = 0$ if C_{Φ} is hyponormal on H^2. We conjecture that the same result is true for A^2. Moreover, the next results are similar to theirs.

Lemma : If $0 < |a| < 1$ or if $|a| = 1$ and $\Phi'(a) = 1$, then neither C_{Φ} nor C^*_{Φ} is hyponormal on A^2.

Proof : The spectral radius theorem implies $r(C_{\Phi})=1$ but $\|C_{\Phi}\| > 1$. Therefore, [4,p.141] implies that neither C_{Φ} nor C^*_{Φ} is hyponormal on A^2.

We note that in the lemma neither C_{Φ} nor C^*_{Φ} is subnormal on A^2 since subnormality implies hyponormality[4,p. 140].

Theorem 4.2 : If C^*_{Φ} is hyponormal on A^2, then $|a| = 1$ and $\Phi'(a) < 1$, or else C_{Φ} is normal on A^2.

Proof : By the lemma we need only examine the case $\Phi(0) = 0$.
We have $S = z^k A^2$ is an invariant subspace of C_{Φ} on

A^2 $\forall$ positive integers k. Hence , $S^{\perp}$ is an invariant subspace of C^*_{Φ} .Sine $S^{\perp}$ is finite dimensional and C^*_{Φ} hyponormal on it, [4,p. 142] implies that C^*_{Φ} is normal on $S^{\perp}$.
Therefore, by [1] $\Phi(z) = \alpha z$ for some α with $|\alpha| \leq 1$ and consequently C_{Φ} is normal.

5. Example . Let $\Phi(z) = z^m$, $m = 2,3,\ldots$ and C_{Φ} be the induced composition operator on A^2. We prove that

a) $\sigma(C_{\Phi}) = \{\lambda \in C : |\lambda| \leq 1/\sqrt{m}\} \cup \{1\}$ (5.1)
Where $\sigma(C_{\Phi})$ is the spectrum of C_{Φ}.

b) C_{Φ} is bounded below by $1/\sqrt{m}$

c) C_{Φ} is not subnormal.

Proof : a) Let $f(z) = \Sigma^{\infty}_{k=0} a_k z^k$ and $g(z) \neq bz$ where $b = 0$.
Suppose $\lambda \neq 1$ and $(C_{\Phi} - \lambda I)(f) = g$. Then

$$\sum_{k-0}^{\infty} a_k z^{mk} = bz + \sum_{j-0}^{\infty} \lambda a_j z^j$$

Fixing m and equating the corresponding coefficients we get

$$a_1 = -\frac{b}{\lambda} \quad \text{and} \quad a_{mk} = \frac{a_k}{\lambda}, \quad k=1,2,3,\ldots$$

Hence,

$$a_{m^n} = -\frac{b}{\lambda^{n+1}}, \; n=1,2,3,\ldots$$

Thus,

$$\|f\|^2 = \sum_{k=0}^{\infty} \frac{|a_k|^2}{k+1} \geq \sum_{n=1}^{\infty} \frac{|a_{m^n}|^2}{m^n+1} = |b|^2 \sum_{n=1}^{\infty} \frac{1}{\lambda^{2n+2}(m^n+1)}$$

Therefore, the ratio test implies there does not exist $f \in A^2$ such that $(C_\Phi - \lambda I)(f) = g$ if $|\lambda| < 1/\sqrt{m}$ which means

$$\{\lambda \in \mathbb{C} : |\lambda| < 1/\sqrt{m}\} \subseteq \sigma(C_\Phi) \qquad (5.2)$$

Next let $A^2_0 = \{ f \in A^2 : f(0) = 0 \}$. If $C_\Phi|_{A^2_0}$ is the restriction of C_Φ to A^2_0 and $f(z) = \Sigma^{\infty}_{k=1} a_k z^k \in A^2$, then for each $n = 1,2,\ldots$ we have

$$\|(C_\Phi|_{A^2_0})^n(f)\|^2 = \|f \circ \Phi_n\|^2 = \|\sum_{k=1}^{\infty} a_k z^{m^n k}\|^2$$

$$= \sum_{k=1}^{\infty} \frac{|a_k|^2}{(k+1)} \frac{(k+1)}{(m^n k+1)}$$

Since $(k+1)/(m^n k+1)$ decreases to $1/m^n$ as $k \to \infty$ we get

$$(1/m^n)\|f\|^2 \le (C_\phi|_{A_0^2})^n(f)\|^2 \le (2/(m^n+1))\|f\|^2$$

Therefore,

$$(1/\sqrt{m}) \le \|(C_\phi|_{A_0^2})^n\|^{1/n} \le (2/m^n+1)^{1/2n}$$

letting $n \to \infty$ it follows that

$$r(C_\phi|_{A_0^2}) = 1/\sqrt{m} \qquad (5.3)$$

Next if $C_\Phi|_C$ is the restriction of C_Φ to the complex numbers then by [5] the only non-zero solutions of $(C_\Phi - \lambda I)(f) = 0$ is $\lambda = 1$ and f constant. So if $\lambda \neq 1$, then the kernel of $C_\Phi - \lambda I$ is zero. Moreover, $\forall$ constant α

$$(C_\phi|_C - \lambda I)\left(\frac{\alpha}{1-\lambda}\right) = \alpha$$

i.e., $C_\Phi|_C - \lambda I$ is onto and hence invertible. Therefore,

$\sigma(C_\Phi|_C) = \{1\}$.

Finally , since

$$\sigma(C_\phi) = \sigma(C_\phi|_{A_0^2 + C_\phi|_C}) = \sigma(C_\phi|_{A_0^2}) \bigcup \sigma(C_\phi|_C)$$

(see e.g., [4,p. 43]) and observing that 1 is an eigenvalue of C_Φ (5.2) and (5.3) imply (5.1).

(b) let $f(z) = \Sigma^\infty_{K=0} a_k z^k \in A^2$. Since $(k+1)/(mk+1)$ decreases to $1/m$ as $k \to \infty$ we see that C_Φ is bounded below by $1/\sqrt{m}$ from

$$\|C_\phi f\|^2 = \|\sum_{k=0}^\infty a_k z^{mk}\|^2 = \sum_{k=0}^\infty \frac{|a_k|^2}{(k+1)} \left(\frac{k+1}{mk+1}\right) \geq \frac{1}{m}\|f\|^2$$

(c) Theorem 4.1 implies that C_Φ is not subnormal because

$$\|C_\phi^2 e_k\| = \sqrt{\frac{2}{m^2+1}} < \frac{2}{m+1} = \|C_\phi e_1\|^2$$

We close this example by pointing out that

$$C_\phi e_k = \sqrt{\frac{1+k}{1+mk}}\, e_{mk}\ ,\ k=0,1,2,\ldots$$

and

$$C_\phi^* e_k = \begin{cases} \sqrt{\dfrac{1+(k/m)}{1+k}}\, e_{k/m} & if\ (k/m) \in \mathbb{N} \\ 0 & if\ (k/m) \notin \mathbb{N} \end{cases}$$

Where N is the natural numbers.

REFERENCES

1. D.M. Boyd, Composition Operators on the Bergman Space, Collog. Math., 34 (1975), 127-136.
2. R. K. Campbell-Wright, On the Equivalence of Composition Operators, thesis, Purdue University, 1989.
3. C. Caratheodory, Theory of functions of a Complex Variable, vol. II, second English edition, Chelsea Publishing Co., New York 1960.
4. John B. Conway, Subnormal Operators, Pitman Publishing Inc., Boston, 1981.
5. Carl C. Cowen, Iteration and the Solution of Functional Equations for Functions Analytic in the Unit Disk, Trans Amer. Math. Soc. 256 (1981),69-95(MR 82 i # 30036)
6. _____ , Composition Operators on H^2,J. Operator Theory 9(1983), 77-106(MR 84d # 47038).
7. C. C. Cowen and T. L. Kriete, Subnormality and Composition Operators on H^2, J. Funct. Anal. 81(1988), 298-319.
8. A. Denjoy, Sur l'iteration des fonctions analytiques, C. R. Acad. Sci. Paris 182(1926), 255-275.
9. J. Wollf, Sur l'iteration des fonctions, Cn R. Acad. Sci. Paris 182(1926) 42-43, 200-201.